

Title: Permeation of chlorhexidine from alcoholic and aqueous solutions within excised human skin

Short running title: Skin penetration of chlorhexidine

T. J. Karpanen^{1*}, T. Worthington¹, B. R. Conway¹, A. C. Hilton¹, T. S. J. Elliott² and P. A. Lambert¹.

1. Life & Health Sciences, Aston University, Aston Triangle, Birmingham B4 7ET, UK

2. Selly Oak Hospital, University Hospital Birmingham NHS Foundation Trust, Raddlebarn Road, Selly Oak, Birmingham B29 6JD.

* Corresponding author. Mailing address: Life and Health Sciences, Aston University, Aston Triangle, B4 7ET, UK. Tel. (44) 121 204 3951, fax. (44) 121 204 4187. E-mail: karpan@aston.ac.uk.

Key words

Skin permeation, antiseptic, HPLC, Franz diffusion cell, chlorhexidine

19 **Dear Editor,**

20
21 **Chlorhexidine** digluconate (CHG) is widely used in the clinical setting for skin
22 antiseptics prior to incision or insertion of medical devices for example central venous
23 catheters (11-13), however, its permeation into skin is limited (6, 7, 9, 10, 16). In a recent
24 study, we demonstrated the limited penetration of CHG in a skin model comprising full
25 thickness excised human skin following application of 2% (w/v) aqueous CHG (9). The
26 aim of this current study was to compare the penetration of chlorhexidine into skin
27 following topical application of 2% (w/v) CHG in 70% (v/v) iso-propyl alcohol (IPA)
28 compared to aqueous CHG.

29 Skin permeation studies were performed on full-thickness excised human skin
30 as described previously (9). Many differences between animal models and human skin
31 absorption have been shown to be permeant-specific, and due to the applied nature of this
32 work, it was considered that human skin must be used (3, 5). Due to the limitation with
33 availability of fresh human skin, frozen skin is most often used as a model to minimize
34 interpersonal variability on the donor skin permeability. Human skin has benefits over
35 animal skin and storage of human skin for a prolonged times has been shown not to have
36 a significant effect on skin permeability (3, 5). In brief, excised human skin was exposed
37 to aqueous and alcoholic CHG for clinically relevant time periods of 2 min and 30 min in
38 a Franz cell diffusion model. The concentration of CHG in serial skin sections was
39 determined by high-performance liquid chromatography.

40 Overall, following a 2 and 30 min exposure, skin penetration of CHG from both
41 aqueous and alcoholic solutions was limited (Figures 1 and 2). At skin depths of ≥ 300

42 μm the concentration of CHG detected from both solutions was negligible ($<0.0008 \mu\text{g}/$
43 mg tissue). In addition, CHG did not completely permeate through the full thickness
44 human skin model.

45 The concentration of CHG recovered within the top $100 \mu\text{m}$ skin sections was
46 significantly less following a 2 min exposure to alcoholic CHG than that following
47 similar exposure to aqueous CHG [mean CHG concentration (\pm s.e.) of $0.023 (\pm 0.007)$
48 μg and $0.157 (\pm 0.047) \mu\text{g}$ per mg tissue for CHG/IPA and CHG respectively, ($p=$
49 0.008)]. Following a 30 min exposure, there was no significant difference in skin
50 penetration of CHG from alcoholic and aqueous solutions within the model ($p>0.05$).

51 The results from this study clearly demonstrate the limited permeation of CHG
52 within a human skin model following application of either alcoholic or aqueous solutions.
53 Moreover, the negligible concentrations of CHG detected at skin depths of $>300 \mu\text{m}$ may
54 indeed allow for microorganisms residing in the deeper layers, for example around hair
55 follicles, to survive the skin antisepsis procedures recommended in the current EPIC
56 guidelines (13).

57 Whilst chlorhexidine in alcoholic solution has clearly been shown to have
58 superior antimicrobial activity compared to aqueous CHG (1, 8), their efficacy in
59 reducing catheter colonization and infection is comparable (14). Alcohol, at a
60 concentration of 70% (v/v) has rapid antimicrobial activity against a broad spectrum of
61 microorganisms (2). However, it has also been shown to extract important lipid
62 components of the stratum corneum (SC) and to cause dehydration of SC proteins, thus
63 potentially compromising the permeation of CHG within the skin (4, 15). These results

64 clearly lay the foundation for further research within the field of skin antisepsis with a
65 view to developing improved formulation strategies for use of CHG in clinical practice.

ACCEPTED

66 **1. Adams, D., M. Quayum, T. Worthington, P. Lambert, and T. Elliott.** 2005.
67 Evaluation of a 2% chlorhexidine gluconate in 70% isopropyl alcohol skin disinfectant. J
68 Hosp Infect **61**:287-90.

69 **2. Adams, D. H.** 2006. An evaluation of three strategies to reduce device related
70 infections associated with hypodermic needles and peripheral vascular catheters. Aston
71 University, Birmingham.

72 **3. Bronaugh R. L., R. F. Stewart, and M. Simon.** 1986. Methods for in vitro
73 percutaneous absorption studies. VII: Use of excised human skin. J Pharm Sci 75: 1094-
74 1097.

75 **4. dos Anjos, J. L., D. de Sousa Neto, and A. Alonso.** 2007. Effects of ethanol/l-
76 menthol on the dynamics and partitioning of spin-labeled lipids in the stratum corneum.
77 Eur J Pharm Biopharm **67**:406-12.

78 **5. Harrison, S.M., Barry, B.W. & Dugard, P.H.** 1984. Effects of freezing on human
79 skin permeability. J Pharm Pharmacol 36: 261-262.

80 **6. Hendley, J. O., and K. M. Ashe.** 1991. Effect of Topical Antimicrobial Treatment on
81 Aerobic Bacteria in the Stratum Corneum of Human Skin. Antimicrobial Agents and
82 Chemotherapy **35**:627-631.

83 **7. Hendley, J. O., and K. M. Ashe.** 2003. Eradication of Resident Bacteria of Normal
84 Human Skin by Antimicrobial Ointment. Antimicrobial Agents and Chemotherapy
85 **47**:1988-1990.

86 **8. Hibbard, J. S., G. K. Mulberry, and A. R. Brady.** 2002. A clinical study comparing
87 the skin antisepsis and safety of Chloraprep, 70% isopropyl alcohol, and 2% aqueous
88 chlorhexidine. J Infus Nurs **25**:244-9.

- 89 **9. Karpanen, T. J., T. Worthington, B. R. Conway, A. C. Hilton, T. S. Elliott, and P.**
90 **A. Lambert.** 2008. Penetration of chlorhexidine into human skin. *Antimicrob Agents*
91 *Chemother* **52**:3633-3636.
- 92 **10. Lafforgue, C., L. Carret, F. Falson, M. E. Reverdy, and J. Freney.** 1997.
93 Percutaneous absorption of a chlorhexidine digluconate solution. *International Journal of*
94 *Pharmaceutics* **147**:243-246.
- 95 **11. McDonnell, G., and A. D. Russell.** 1999. Antiseptics and disinfectants: activity,
96 action, and resistance. *Clin Microbiol Rev* **12**:147-79.
- 97 **12. O'Grady, N. P., M. Alexander, E. P. Dellinger, J. L. Gerberding, S. O. Heard, D.**
98 **G. Maki, H. Masur, R. D. McCormick, L. A. Mermel, M. L. Pearson, Raad, II, A.**
99 **Randolph, and R. A. Weinstein.** 2002. Guidelines for the prevention of intravascular
100 catheter-related infections. Centers for Disease Control and Prevention. *MMWR*
101 *Recomm Rep* **51**:1-29.
- 102 **13. Pratt, R. J., C. M. Pellowe, J. A. Wilson, H. P. Loveday, P. J. Harper, S. R.**
103 **Jones, C. McDougall, and M. H. Wilcox.** 2007. epic2: National evidence-based
104 guidelines for preventing healthcare-associated infections in NHS hospitals in England. *J*
105 *Hosp Infect* **65 Suppl 1**:S1-64.
- 106 **14. Valles, J., I. Fernandez, D. Alcaraz, E. Chacon, A. Cazorla, M. Canals, D.**
107 **Mariscal, D. Fontanals, and A. Moron.** 2008. Prospective Randomized Trial of 3
108 Antiseptic Solutions for Prevention of Catheter Colonization in an Intensive Care Unit for
109 Adult Patients. *Infect Control Hosp Epidemiol* **29**:847-53.

- 110 **15. Van der Merwe, D., and J. E. Riviere.** 2005. Comparative studies on the effects of
111 water, ethanol and water/ethanol mixtures on chemical partitioning into porcine stratum
112 corneum and silastic membrane. *Toxicol In Vitro* **19**:69-77.
- 113 **16. Wang, J. C., R. R. Williams, L. Wang, and J. Loder.** 1990. In vitro skin
114 permeation and bioassay of chlorhexidine phosphanilate, a new antimicrobial agent.
115 *Pharm Res* **7**:995-1002.




Figure 1. Penetration profile showing the location and concentration of chlorhexidine ($\mu\text{g}/\text{mg}$ tissue) in excised human skin after 2 min ($n=15$) exposure to 2% (w/v) chlorhexidine digluconate in 70% (v/v) isopropyl alcohol and aqueous 2% (w/v) CHG (mean \pm s.e., * $p=0.003$).




Figure 2. Penetration profile showing the location and concentration of chlorhexidine ($\mu\text{g}/\text{mg}$ tissue) in excised human skin after 30 min ($n=15$) exposure to 2% (w/v) chlorhexidine digluconate in 70% (v/v) isopropyl alcohol and aqueous 2% (w/v) CHG (mean \pm s.e).